

LEHMER-TYPE CONGRUENCES FOR LACUNARY HARMONIC SUMS MODULO p^2

HAO PAN

ABSTRACT. In this paper, we establish some Lehmer-type congruences for lacunary harmonic sums modulo p^2 .

1. INTRODUCTION

The well-known Wolstenholme's harmonic series congruence asserts that

$$\sum_{k=1}^{p-1} \frac{1}{k} \equiv 0 \pmod{p^2} \quad (1.1)$$

for each prime $p \geq 5$. With help of (1.1), Wolstenholme [9] proved that

$$\binom{mp}{np} \equiv \binom{m}{n} \pmod{p^3}.$$

for any $m, n \geq 1$ and prime $p \geq 5$. In 1938, Lehmer [2] discovered an interesting congruence as follows:

$$\sum_{j=1}^{(p-1)/2} \frac{1}{j} \equiv -\frac{2^p - 2}{p} + \frac{(2^{p-1} - 1)^2}{p} \pmod{p^2}$$

for each prime $p \geq 3$.

Define

$$\mathcal{H}_{r,m}(n) = \sum_{\substack{1 \leq k \leq n \\ k \equiv r \pmod{m}}} \frac{1}{k}.$$

Clearly, with help of 1.1, Lehmer's congruence can be rewritten as

$$\mathcal{H}_{p,2}(p-1) \equiv \frac{2^{p-1} - 1}{p} - \frac{(2^{p-1} - 1)^2}{2p} \pmod{p}. \quad (1.2)$$

In fact, Lehmer also proved three another congruences in the same flavor:

$$\mathcal{H}_{p,3}(p-1) \equiv \frac{3^{p-1} - 1}{2p} - \frac{(3^{p-1} - 1)^2}{4p} \pmod{p^2}, \quad (1.3)$$

$$\mathcal{H}_{p,4}(p-1) \equiv \frac{3(2^{p-1} - 1)}{4p} - \frac{3(2^{p-1} - 1)^2}{8p} \pmod{p^2} \quad (1.4)$$

2000 *Mathematics Subject Classification.* Primary 11A07; Secondary 11B65.

The author was supported by the National Natural Science Foundation of China (Grant No. 10771135).

and

$$\mathcal{H}_{p,6}(p-1) \equiv \frac{2^{p-1}-1}{3p} + \frac{3^{p-1}-1}{4p} - \frac{(2^{p-1}-1)^2}{6p} - \frac{(3^{p-1}-1)^2}{8p} \pmod{p^2}, \quad (1.5)$$

where $p \geq 5$ is a prime. The proofs of (1.2), (1.3), (1.4) and (1.5) are based on the values of Bernoulli polynomial $B_{p(p-1)}(x)$ at $x = 1/2, 1/3, 1/4, 1/6$.

However, no another congruence for $\mathcal{H}_{p,m}(p-1)$ modulo p^2 is known, partly since very few is known on the values of $B_{p(p-1)}(n/m)$ when $m \neq 1, 2, 3, 4, 6$. Some Lehmer-type congruences modulo p (not modulo p^2 !) have been proved in [8, 3, 4, 5, 6, 7]. In this paper, we shall investigate the Lehmer-type congruences modulo p^2 .

Define

$$\mathcal{T}_{r,m}(n) = \sum_{\substack{0 \leq k \leq n \\ k \equiv r \pmod{m}}} \binom{n}{k} \quad \text{and} \quad \mathcal{T}_{r,m}^*(n) = \sum_{\substack{0 \leq k \leq n \\ k \equiv r \pmod{m}}} (-1)^k \binom{n}{k}.$$

Clearly $\mathcal{T}_{r,m}^*(n) = (-1)^n \mathcal{T}_{n-r,m}^*(n)$ and

$$\mathcal{T}_{r,m}^*(n) = \begin{cases} (-1)^r \mathcal{T}_{r,m}(n) & \text{if } m \text{ is even,} \\ (-1)^r (\mathcal{T}_{r,2m}(n) - \mathcal{T}_{m+r,2m}(n)) & \text{if } m \text{ is odd.} \end{cases}$$

As we shall see soon, it is not difficult to show that

$$\mathcal{H}_{r,m}(p-1) \equiv -\frac{\mathcal{T}_{r,m}^*(p) - \delta_{r,m}(p)}{p} \pmod{p},$$

where

$$\delta_{r,m}(p) = \begin{cases} 1 & \text{if } r \equiv 0 \pmod{m}, \\ -1 & \text{if } r \equiv p \pmod{m}, \\ 0 & \text{otherwise.} \end{cases} \quad (1.6)$$

Theorem 1.1. *Let $m \geq 2$ be an integer and let $p > 3$ be a prime with $p \neq m$. Then*

$$\mathcal{H}_{p,m}(p-1) \equiv -\frac{2\mathcal{T}_{p,m}^*(p) + 2}{p} + \frac{\mathcal{T}_{p,m}^*(2p) + 2}{4p} \pmod{p^2}. \quad (1.7)$$

Let us see how (1.2) follows from Theorem 1.1. Clearly we have $\mathcal{T}_{0,2}^*(n) = 2^{n-1}$ and $\mathcal{T}_{1,2}^*(n) = -2^{n-1}$. Hence in view of (1.7), for any prime $p \geq 5$,

$$\mathcal{H}_{p,2}(p-1) \equiv -\frac{2\mathcal{T}_{p,2}^*(p) + 2}{p} + \frac{\mathcal{T}_{p,2}^*(2p) + 2}{4p} = \frac{2^p - 2}{p} - \frac{2^{2p-1} - 2}{4p} \pmod{p^2}.$$

In [7], Sun had showed that $\mathcal{T}_{r,m}(n)$ can be expressed in terms of some linearly recurrent sequences with orders not exceeding $\phi(m)/2$, where ϕ is the Euler totient function. Thus in view of Theorem 1.1, for each m , we always have a Lehmer-type congruence for $\mathcal{H}_{p,m}(p-1)$ modulo p^2 , involving some linearly recurrent sequences.

However, as we shall see later, (1.7) is not suitable to derive (1.3), (1.4) and (1.5). So we need the following theorem.

Theorem 1.2. *Let $m \geq 2$ be an integer and let $p > 3$ be a prime with $p \neq m$. Then*

$$\mathcal{H}_{p,m}(p-1) \equiv -\frac{\mathcal{T}_{p,m}^*(2p) + 2}{4p} - \frac{p}{2} \sum_{\substack{1 \leq r \leq m \\ 2r \not\equiv p \pmod{m}}} \mathcal{H}_{r,m}(p-1)^2 \pmod{p^2}. \quad (1.8)$$

When $m = 3$, we have $\mathcal{T}_{p,3}^*(2p) = -2 \times 3^{p-1}$ (cf. [3, Theorem 1.9] and [7, Theorem 3.2]). Thus by (1.8), we get

$$\mathcal{H}_{p,3}(p-1) \equiv -\frac{\mathcal{T}_{p,3}^*(2p) + 2}{4p} - p \left(\frac{\mathcal{T}_{p,3}^*(2p) + 2}{4p} \right)^2 = \frac{3^{p-1} - 1}{2p} - \frac{(3^{p-1} - 1)^2}{4p} \pmod{p^2},$$

since

$$\mathcal{H}_{0,3}(p-1) \equiv -\mathcal{H}_{p,3}(p-1) \equiv \frac{\mathcal{T}_{p,3}^*(2p) + 2}{4p} \pmod{p}.$$

Let us apply Theorem 1.2 to obtain more Lehmer's type congruences. The Fibonacci numbers $F_0, F_1, F_2, \dots$ are given by $F_0 = 0$, $F_1 = 1$ and $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$. It is well-known that $F_p \equiv \left(\frac{5}{p}\right) \pmod{p}$ and $F_{p-\left(\frac{5}{p}\right)} \equiv 0 \pmod{p}$ for prime $p \neq 2, 5$, where $\left(\frac{\cdot}{p}\right)$ is the Legendre symbol. Williams [8] proved that

$$\frac{2}{5} \sum_{1 \leq k \leq 4p/5-1} \frac{(-1)^k}{k} \equiv \frac{F_{p-\left(\frac{5}{p}\right)}}{p} \pmod{p}$$

for prime $p \neq 2, 5$. Subsequently Sun and Sun [6, Corollary 3] proved that

$$\mathcal{H}_{2p,5}(p-1) \equiv -\mathcal{H}_{-p,5}(p-1) \equiv -\frac{F_{p-\left(\frac{5}{p}\right)}}{2p} \pmod{p}. \quad (1.9)$$

We have a Lehmer-type congruences as follows.

Theorem 1.3. *Suppose that $p > 5$ is a prime. Then*

$$\mathcal{H}_{p,5}(p-1) \equiv \frac{5^{\frac{p-1}{2}} F_p - 1}{p} - \frac{5^{p-1} F_{2p-\left(\frac{5}{p}\right)} - 1}{4p} \pmod{p^2}. \quad (1.10)$$

The Pell numbers $P_0, P_1, P_2, \dots$ are given by $P_0 = 0$, $P_1 = 1$ and $P_n = 2P_{n-1} + P_{n-2}$ for $n \geq 2$. We know $P_p \equiv \left(\frac{2}{p}\right) \pmod{p}$ and $P_{p-\left(\frac{2}{p}\right)} \equiv 0 \pmod{p}$ for every odd prime p . In [4], Sun proved that

$$(-1)^{\frac{p-1}{2}} \sum_{1 \leq k \leq (p+1)/4} \frac{(-1)^k}{2k-1} \equiv -\frac{1}{4} \sum_{k=1}^{\frac{p-1}{2}} \frac{2^k}{k} \equiv \frac{P_{p-\left(\frac{2}{p}\right)}}{p} \pmod{p}$$

for odd prime p . Similarly, we have a Lehmer-type congruence involving Pell numbers.

Theorem 1.4. *Suppose that $p > 3$ is a prime. Then*

$$\mathcal{H}_{p,8}(p-1) \equiv \frac{2^{2p-4} + 2^{p-3} + 2^{\frac{p-3}{2}} P_p - 1}{p} - \frac{2^{4p-6} + 2^{2p-4} + 2^{p-2} P_{2p-\left(\frac{2}{p}\right)} - 1}{4p} \pmod{p^2}. \quad (1.11)$$

We shall prove Theorems 1.1 and 1.2 in Section 2. And the proofs of Theorems 1.3 and 1.4 will be given in Section 3.

2. PROOF THEOREMS 1.1 AND 1.2

Lemma 2.1. *Suppose that p is a prime. Then*

$$\frac{1}{p} \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} (-a)^k \binom{p}{k} \equiv - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{k} + p \sum_{\substack{1 \leq j < k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{jk} \pmod{p^2} \quad (2.1)$$

and

$$\begin{aligned} \frac{1}{2p} \sum_{\substack{1 \leq k \leq 2p-1, k \neq p \\ k \equiv r \pmod{m}}} (-a)^k \binom{2p}{k} &\equiv - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{k} - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \frac{a^{2p-k}}{k} \\ &\quad + 2p \sum_{\substack{1 \leq j < k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{jk} + 2p \sum_{\substack{1 \leq j < k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \frac{a^{2-k}}{jk} \pmod{p^2}. \end{aligned} \quad (2.2)$$

Proof.

$$\begin{aligned} \frac{1}{p} \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} (-a)^k \binom{p}{k} &= \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{(-a)^k}{k} \prod_{j=1}^{k-1} \left(\frac{p}{j} - 1 \right) \\ &\equiv - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{k} + \sum_{\substack{2 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{k} \sum_{j=1}^{k-1} \frac{p}{j} \pmod{p^2}. \end{aligned}$$

Similarly,

$$\begin{aligned} &\frac{1}{2p} \sum_{\substack{1 \leq k \leq 2p-1, k \neq p \\ k \equiv r \pmod{m}}} (-a)^k \binom{2p}{k} \\ &= \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{(-a)^k}{k} \binom{2p-1}{k-1} + \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \frac{(-a)^{2p-k}}{2p-k} \binom{2p-1}{k}. \end{aligned}$$

We have

$$\sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{(-a)^k}{k} \binom{2p-1}{k-1} \equiv - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{k} + 2p \sum_{\substack{1 \leq j < k \leq p-1 \\ k \equiv r \pmod{m}}} \frac{a^k}{jk} \pmod{p^2}.$$

And

$$\begin{aligned}
 & \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \frac{(-a)^{2p-k}}{2p-k} \binom{2p-1}{k} \\
 & \equiv \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \frac{a^{2p-k}}{2p-k} - 2p \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \frac{a^{2-k}}{2p-k} \sum_{j=1}^k \frac{1}{j} \\
 & \equiv - \sum_{\substack{1 \leq k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \left(\frac{a^{2p-k}}{k} + 2p \cdot \frac{a^{2-k}}{k^2} \right) + 2p \sum_{\substack{1 \leq j \leq k \leq p-1 \\ k \equiv 2p-r \pmod{m}}} \frac{a^{2-k}}{jk} \pmod{p^2}.
 \end{aligned}$$

We are done. $\square$

Define

$$\mathcal{S}_{r,m}(n) = \sum_{\substack{2 \leq k \leq n \\ k \equiv r \pmod{m}}} \frac{1}{k} \sum_{j=1}^{k-1} \frac{1}{j}.$$

Substituting $a = 1$ in (2.1), we get

Corollary 2.1. *Suppose that $m \geq 2$. Then*

$$\mathcal{H}_{r,m}(p-1) \equiv -\frac{\mathcal{T}_{r,m}^*(p) - \delta_{r,m}(p)}{p} + p\mathcal{S}_{r,m}(p-1) \pmod{p^2}, \quad (2.3)$$

where $\delta_{r,m}(p)$ is same as the one defined in (1.6). In particular,

$$\mathcal{H}_{p,m}(p-1) \equiv -\frac{\mathcal{T}_{p,m}^*(p) + 1}{p} + p\mathcal{S}_{p,m}(p-1) \pmod{p^2}. \quad (2.4)$$

Substituting $r = p, p + m/2$ and $a = 1$ in (2.2) and noting that $\binom{2p}{p} \equiv 2 \pmod{p^3}$, we have

Corollary 2.2. *Suppose that $m \geq 2$. Then*

$$\mathcal{H}_{p,m}(p-1) \equiv -\frac{\mathcal{T}_{p,m}^*(2p) + 2}{4p} + 2p\mathcal{S}_{p,m}(p-1) \pmod{p^2}. \quad (2.5)$$

And if m is even, then

$$\mathcal{H}_{p+m/2,m}(p-1) \equiv -\frac{\mathcal{T}_{p+m/2,m}^*(2p)}{4p} + 2p\mathcal{S}_{p+m/2,m}(p-1) \pmod{p^2}. \quad (2.6)$$

Combining (2.4) and (2.5), we get

$$p\mathcal{S}_{p,m}(p-1) \equiv -\frac{\mathcal{T}_{p,m}^*(p) + 1}{p} + \frac{\mathcal{T}_{p,m}^*(2p) + 2}{4p} \pmod{p^2},$$

and Theorem 1.1 easily follows.

Lemma 2.2.

$$\sum_{r=1}^m \mathcal{T}_{r,m}^*(n) \mathcal{T}_{r+s,m}^*(n) = (-1)^n \mathcal{T}_{n+s,m}^*(2n).$$

Proof. Let ζ be a primitive m -th root of unity. Clearly,

$$\mathcal{T}_{r,m}^*(n) = \frac{1}{m} \sum_{k=0}^n (-1)^k \binom{n}{k} \sum_{t=1}^m \zeta^{(k-r)t} = \frac{1}{m} \sum_{t=1}^m \zeta^{-rt} (1 - \zeta^t)^n.$$

Hence

$$\begin{aligned} \sum_{r=1}^m \mathcal{T}_{r,m}^*(n) \mathcal{T}_{r+s,m}^*(n) &= \sum_{r=1}^m \frac{1}{m^2} \sum_{1 \leq t_1, t_2 \leq m} \zeta^{-r(t_1+t_2)-st_2} (1 - \zeta^{t_1})^n (1 - \zeta^{t_2})^n \\ &= \frac{(-1)^n}{m} \sum_{t=1}^m \zeta^{-(n+s)t} (1 - \zeta^t)^{2n} = (-1)^n \mathcal{T}_{n+s,m}^*(2n). \end{aligned}$$

□

By Lemma 2.2, we have

$$\mathcal{T}_{p,m}^*(2p) = - \sum_{r=1}^m \mathcal{T}_{r,m}^*(p)^2.$$

Hence

$$\begin{aligned} \mathcal{S}_{p,m}(p-1) &\equiv \frac{\mathcal{T}_{0,m}^*(p) - 1}{2p^2} - \frac{\mathcal{T}_{p,m}^*(p) + 1}{2p^2} - \frac{\sum_{r=1}^m \mathcal{T}_{r,m}^*(p)^2 - 2}{4p^2} \\ &= - \sum_{\substack{1 \leq r \leq m \\ r \not\equiv 0, p \pmod{m}}} \frac{\mathcal{T}_{r,m}^*(p)^2}{4p^2} - \frac{(\mathcal{T}_{0,m}^*(p) - 1)^2}{4p^2} - \frac{(\mathcal{T}_{p,m}^*(p) + 1)^2}{4p^2} \\ &\equiv - \frac{1}{4} \sum_{r=1}^m \mathcal{H}_{r,m}(p-1)^2 \pmod{p}. \end{aligned} \tag{2.7}$$

And since $\mathcal{H}_{p-r,m}(p-1) \equiv -\mathcal{H}_{r,m}(p-1) \pmod{p}$, $\mathcal{H}_{r,m}(p-1) \equiv 0 \pmod{p}$ provided that $2r \equiv p \pmod{m}$. So we also have

$$\mathcal{S}_{p,m}(p-1) \equiv -\frac{1}{4} \sum_{\substack{1 \leq r \leq m \\ 2r \not\equiv p \pmod{m}}} \mathcal{H}_{r,m}(p-1)^2 \pmod{p}. \tag{2.8}$$

Thus by (2.5), Theorem 1.2 is concluded.

3. FERMAT'S QUOTIENT AND PELL'S QUOTIENT

Let L_n be the Lucas numbers given by $L_0 = 2$, $L_1 = 1$ and $L_n = L_{n-1} + L_{n-2}$ for $n \geq 2$. We require the following result of Sun and Sun on $\mathcal{T}_{r,10}(n)$.

Lemma 3.1. [6, Theorem 1] *Let n be a positive odd integer. If $n \equiv 1 \pmod{4}$, then*

$$\begin{aligned} 10\mathcal{T}_{\frac{n-1}{2},10}(n) &= 2^n + L_{n+1} + 5^{\frac{n+3}{4}} F_{\frac{n+1}{2}}, & 10\mathcal{T}_{\frac{n+3}{2},10}(n) &= 2^n - L_{n-1} + 5^{\frac{n+3}{4}} F_{\frac{n-1}{2}}, \\ 10\mathcal{T}_{\frac{n+7}{2},10}(n) &= 2^n - L_{n-1} - 5^{\frac{n+3}{4}} F_{\frac{n-1}{2}}, & 10\mathcal{T}_{\frac{n+11}{2},10}(n) &= 2^n + L_{n+1} - 5^{\frac{n+3}{4}} F_{\frac{n+1}{2}}. \end{aligned}$$

And if $n \equiv 3 \pmod{4}$, then

$$\begin{aligned} 10\mathcal{T}_{\frac{n-1}{2},10}(n) &= 2^n + L_{n+1} + 5^{\frac{n+1}{4}} L_{\frac{n+1}{2}}, & 10\mathcal{T}_{\frac{n+3}{2},10}(n) &= 2^n - L_{n-1} + 5^{\frac{n+1}{4}} L_{\frac{n-1}{2}}, \\ 10\mathcal{T}_{\frac{n+7}{2},10}(n) &= 2^n - L_{n-1} - 5^{\frac{n+1}{4}} L_{\frac{n-1}{2}}, & 10\mathcal{T}_{\frac{n+11}{2},10}(n) &= 2^n + L_{n+1} - 5^{\frac{n+1}{4}} L_{\frac{n+1}{2}}. \end{aligned}$$

Furthermore, for every odd n ,

$$10\mathcal{T}_{\frac{n+13}{2},10}(n) = 2^n - 2L_n.$$

For each odd $n \geq 1$, since

$$\mathcal{T}_{n,m}^*(2n) = \mathcal{T}_{n,m}^*(2n-1) - \mathcal{T}_{n-1,m}^*(2n-1) = -2\mathcal{T}_{n-1,m}^*(2n-1)$$

and

$$\mathcal{T}_{n+m,2m}^*(2n) = \mathcal{T}_{n+m,2m}^*(2n-1) - \mathcal{T}_{n+m-1,m}^*(2n-1) = -2\mathcal{T}_{n+m-1,m}^*(2n-1),$$

by Lemma 3.1, we get

$$\mathcal{T}_{n,5}^*(2n) = -2 \cdot 5^{\frac{n-1}{2}} F_n. \quad (3.1)$$

Let $p > 5$ be a prime. By (1.8),

$$\mathcal{H}_{p,5}(p-1) \equiv -\frac{\mathcal{T}_{p,5}^*(2p) + 2}{4p} - p(H_{p,5}(p-1)^2 + H_{2p,5}(p-1)^2) \pmod{p^2}.$$

By (1.9), we have

$$\begin{aligned} \mathcal{H}_{p,5}(p-1) &\equiv \frac{5^{\frac{p-1}{2}} F_p - 1}{2p} - p \left(\left(\frac{F_{p-(\frac{5}{p})}}{2p} \right)^2 + \left(\frac{5^{\frac{p-1}{2}} F_p - 1}{2p} \right)^2 \right) \\ &\equiv \frac{5^{\frac{p-1}{2}} F_p - 1}{p} - \frac{5^{p-1} (F_{p-(\frac{5}{p})}^2 + F_p^2) - 1}{4p} \\ &= \frac{5^{\frac{p-1}{2}} F_p - 1}{p} - \frac{5^{p-1} F_{2p-(\frac{5}{p})} - 1}{4p} \pmod{p^2}, \end{aligned}$$

where in the last step we use the fact $F_{2n-1} = F_n^2 + F_{n-1}^2$. Thus the proof of Theorem 1.3 is complete.

Remark. Similarly, we can get

$$\begin{aligned} &\sum_{\substack{1 \leq k \leq p-1 \\ k \equiv p \pmod{5}}} \frac{(-1)^k}{k} \\ &\equiv \frac{5(2^{4p-1} - 2^{2p+3}) + 12L_{4p} + L_{4p-4(\frac{5}{p})} - 112L_{2p} - 4L_{2p-2(\frac{5}{p})} + 378}{400p} \pmod{p^2}. \end{aligned} \quad (3.2)$$

Let Q_n be the Pell-Lucas numbers given by $Q_0 = 2$, $Q_1 = 2$ and $Q_n = 2Q_{n-1} + Q_{n-2}$ for $n \geq 2$. For $\mathcal{T}_{r,8}(n)$, Sun had proved that

Lemma 3.2. [4, Theorem 2.2] *Let n be a positive odd integer. If $n \equiv 1 \pmod{4}$, then*

$$\begin{aligned} 8\mathcal{T}_{\frac{n-1}{2},8}(n) &= 2^n + 2^{\frac{n+1}{2}} + 2^{\frac{n+7}{4}} P_{\frac{n+1}{2}}, & 8\mathcal{T}_{\frac{n+3}{2},8}(n) &= 2^n - 2^{\frac{n+1}{2}} + 2^{\frac{n+7}{4}} P_{\frac{n-1}{2}}, \\ 8\mathcal{T}_{\frac{n+7}{2},8}(n) &= 2^n - 2^{\frac{n+1}{2}} - 2^{\frac{n+7}{4}} P_{\frac{n-1}{2}}, & 8\mathcal{T}_{\frac{n+11}{2},8}(n) &= 2^n + 2^{\frac{n+1}{2}} - 2^{\frac{n+7}{4}} P_{\frac{n+1}{2}}. \end{aligned}$$

And if $n \equiv 3 \pmod{4}$, then

$$\begin{aligned} 8\mathcal{T}_{\frac{n-1}{2},8}(n) &= 2^n + 2^{\frac{n+1}{2}} + 2^{\frac{n+1}{4}} Q_{\frac{n+1}{2}}, & 8\mathcal{T}_{\frac{n+3}{2},8}(n) &= 2^n - 2^{\frac{n+1}{2}} + 2^{\frac{n+1}{4}} Q_{\frac{n-1}{2}}, \\ 8\mathcal{T}_{\frac{n+7}{2},8}(n) &= 2^n - 2^{\frac{n+1}{2}} - 2^{\frac{n+1}{4}} Q_{\frac{n-1}{2}}, & 8\mathcal{T}_{\frac{n+11}{2},8}(n) &= 2^n + 2^{\frac{n+1}{2}} - 2^{\frac{n+1}{4}} Q_{\frac{n+1}{2}}. \end{aligned}$$

Thus we have

$$T_{n,8}^*(2n) = -2^{2n-3} - 2^{n-2} - 2^{\frac{n-1}{2}} P_n \quad (3.3)$$

and

$$T_{n+4,8}^*(2n) = -2^{2n-3} - 2^{n-2} + 2^{\frac{n-1}{2}} P_n \quad (3.4)$$

for odd $n \geq 1$. Applying (1.8),

$$\mathcal{H}_{p,8}(p-1) \equiv -\frac{\mathcal{T}_{p,8}^*(2p) + 2}{4p} - p \sum_{0 \leq j \leq 3} \mathcal{H}_{p+2j,8}(p)^2 \pmod{p^2}.$$

By (2.5) and (2.6), we have

$$\mathcal{H}_{p,8}(p-1) \equiv -\frac{\mathcal{T}_{p,8}^*(2p) + 2}{4p} \pmod{p}$$

and

$$\mathcal{H}_{p+4,8}(p-1) \equiv -\frac{\mathcal{T}_{p+4,8}^*(2p)}{4p} \pmod{p}.$$

And in view of (2.3) and Lemma 3.2.

$$\begin{aligned} & \mathcal{H}_{p+2,8}(p-1)^2 + \mathcal{H}_{p+6,8}(p-1)^2 \\ & \equiv \begin{cases} \sum_{i=0}^1 p^{-2} (2^{p-3} - (\frac{2}{p}) 2^{\frac{p-5}{2}} + (-1)^i 2^{\frac{p-5}{4}} P_{(p-(\frac{2}{p})) / 2})^2 \pmod{p} & \text{if } p \equiv 1 \pmod{4}, \\ \sum_{i=0}^1 p^{-2} (2^{p-3} - (\frac{2}{p}) 2^{\frac{p-5}{2}} + (-1)^i 2^{\frac{p-11}{4}} Q_{(p-(\frac{2}{p})) / 2})^2 \pmod{p} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned} \quad (3.5)$$

Lemma 3.3.

$$\begin{aligned} P_{\frac{p-1}{2}} &\equiv 0 \pmod{p}, & P_{\frac{p+1}{2}} &\equiv (-1)^{\frac{p-1}{8}} 2^{\frac{p-1}{4}} \pmod{p}, & \text{if } p &\equiv 1 \pmod{8}, \\ P_{\frac{p-1}{2}} &\equiv (-1)^{\frac{p-3}{8}} 2^{\frac{p-3}{4}} \pmod{p}, & P_{\frac{p+1}{2}} &\equiv (-1)^{\frac{p+5}{8}} 2^{\frac{p-3}{4}} \pmod{p}, & \text{if } p &\equiv 3 \pmod{8}, \\ P_{\frac{p-1}{2}} &\equiv (-1)^{\frac{p-5}{8}} 2^{\frac{p-1}{4}} \pmod{p}, & P_{\frac{p+1}{2}} &\equiv 0 \pmod{p}, & \text{if } p &\equiv 5 \pmod{8}, \\ P_{\frac{p-1}{2}} &\equiv (-1)^{\frac{p+1}{8}} 2^{\frac{p-3}{4}} \pmod{p}, & P_{\frac{p+1}{2}} &\equiv (-1)^{\frac{p+1}{8}} 2^{\frac{p-3}{4}} \pmod{p}, & \text{if } p &\equiv 7 \pmod{8}, \end{aligned} \quad (3.6)$$

and

$$\begin{aligned} Q_{\frac{p-1}{2}} &\equiv (-1)^{\frac{p-1}{8}} 2^{\frac{p+3}{4}} \pmod{p}, & Q_{\frac{p+1}{2}} &\equiv (-1)^{\frac{p-1}{8}} 2^{\frac{p+3}{4}} \pmod{p}, & \text{if } p &\equiv 1 \pmod{8}, \\ Q_{\frac{p-1}{2}} &\equiv (-1)^{\frac{p+5}{8}} 2^{\frac{p+5}{4}} \pmod{p}, & Q_{\frac{p+1}{2}} &\equiv 0 \pmod{p}, & \text{if } p &\equiv 3 \pmod{8}, \\ Q_{\frac{p-1}{2}} &\equiv (-1)^{\frac{p+3}{8}} 2^{\frac{p+3}{4}} \pmod{p}, & Q_{\frac{p+1}{2}} &\equiv (-1)^{\frac{p-5}{8}} 2^{\frac{p+3}{4}} \pmod{p}, & \text{if } p &\equiv 5 \pmod{8}, \\ Q_{\frac{p-1}{2}} &\equiv 0 \pmod{p}, & Q_{\frac{p+1}{2}} &\equiv (-1)^{\frac{p+1}{8}} 2^{\frac{p+1}{4}} \pmod{p}, & \text{if } p &\equiv 7 \pmod{8}. \end{aligned} \quad (3.7)$$

Proof. The congruences in (3.6) were obtained by Sun [4, Theorem 2.3]. And the congruences in (3.7) follows from (3.6), by noting that $Q_n = 2P_{n+1} - 2P_n$ and $Q_{n+1} = 2P_{n+1} + 2P_n$. $\square$

Thus since $P_{(p-(\frac{2}{p})) / 2} Q_{(p-(\frac{2}{p})) / 2} = P_{p-(\frac{2}{p})}$, by (3.5), we have

$$\mathcal{H}_{p+2,8}(p-1)^2 + \mathcal{H}_{p+6,8}(p-1)^2 \equiv \frac{2^{p-1}(2^{\frac{p-1}{2}} - (\frac{2}{p}))^2 + P_{p-(\frac{2}{p})}^2}{8p} \pmod{p}.$$

Observe that

$$\frac{2^{p-1} - 1}{p} = \frac{(2^{\frac{p-1}{2}} + (\frac{2}{p}))(2^{\frac{p-1}{2}} - (\frac{2}{p}))}{p} \equiv 2 \left(\frac{2}{p} \right) \frac{2^{\frac{p-1}{2}} - (\frac{2}{p})}{p} \pmod{p}.$$

Hence

$$\begin{aligned} \mathcal{H}_{p,8}(p-1) &\equiv \frac{2^{2p-4} + 2^{p-3} + 2^{\frac{p-3}{2}} P_p - 1}{p} - \frac{(2^{2p-3} + 2^{p-2} + 2^{\frac{p-1}{2}} P_p)^2 - 4}{16p} \\ &\quad - \frac{(2^{2p-3} + 2^{p-2} - 2^{\frac{p-1}{2}} P_p)^2}{16p} - \frac{2^{p-1}(2^{\frac{p-1}{2}} - (\frac{2}{p}))^2 + 2^{p-1} P_{p-(\frac{2}{p})}^2}{8p} \\ &\equiv \frac{2^{2p-4} + 2^{p-3} + 2^{\frac{p-3}{2}} P_p - 1}{p} - \frac{2^{4p-6} + 2^{2p-4} + 2^{p-2} P_{2p-(\frac{2}{p})} - 1}{4p} \pmod{p^2}, \end{aligned}$$

by noting that $P_{p-(\frac{2}{p})}^2 + P_p^2 = P_{2p-(\frac{2}{p})}$. This concludes the proof of Theorem 1.4. $\square$

Remark. The Bernoulli polynomials $B_n(x)$ are given by

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{B_n(x)}{n!} t^n.$$

In particular, define the Bernoulli number $B_n = B_n(0)$. Granville and Sun [1] proved that

$$B_{p-1}(\{p\}_5/5) - B_{p-1} \equiv \frac{5}{4p} F_{p-(\frac{5}{p})} + \frac{5^p - 5}{4p} \pmod{p}$$

and

$$B_{p-1}(\{p\}_8/8) - B_{p-1} \equiv \frac{2}{p} P_{p-(\frac{2}{p})} + \frac{2^{p+1} - 4}{p} \pmod{p}$$

for prime $p \neq 2, 5$, where $\{p\}_m$ denotes the least non-negative residue of p modulo m . In [5, Theorem 3.3], Sun also proved that

$$m \mathcal{H}_{p,m}(p-1) \equiv \frac{B_{2p-2}(\{p\}_m/m) - B_{2p-2}}{2p-2} - 2 \frac{B_{p-1}(\{p\}_m/m) - B_{p-1}}{p-1} \pmod{p^2}.$$

Now using Theorems 1.3 and 1.4 it is easy to deduce that

$$\frac{B_{p(p-1)}(\{p\}_5/5) - B_{p(p-1)}}{5p(p-1)} \equiv -\frac{5^{\frac{p-1}{2}} F_p - 1}{p} + \frac{5^{p-1} F_{2p-(\frac{5}{p})} - 1}{4p} \pmod{p^2} \quad (3.8)$$

for prime $p > 5$, and

$$\begin{aligned} & \frac{B_{p(p-1)}(\{p\}_8/8) - B_{p(p-1)}}{8p(p-1)} \\ & \equiv -\frac{2^{2p-4} + 2^{p-3} + 2^{\frac{p-3}{2}}P_p - 1}{p} + \frac{2^{4p-6} + 2^{2p-4} + 2^{p-2}P_{2p-(\frac{2}{p})} - 1}{4p} \pmod{p^2} \end{aligned} \quad (3.9)$$

for prime $p > 3$.

Acknowledgment. The author is grateful to Professor Zhi-Wei Sun for his helpful discussions on this paper.

REFERENCES

- [1] A. Granville and Z.-W. Sun, *Values of Bernoulli polynomials*, Pacific J. Math., **172**(1996), 117–137.
- [2] E. Lehmer, *On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson*, Ann. of Math. (2), **39**(1938), 350–360.
- [3] Z.-H. Sun, *The combinatorial sum $\sum_{k=0, k \equiv r \pmod{m}}^n \binom{n}{k}$ and its applications in number theory. I (Chinese)*, J. Nanjing Univ. Math. Biquarterly, **8**(1992), 227–240.
- [4] Z.-H. Sun, *The combinatorial sum $\sum_{k=0, k \equiv r \pmod{m}}^n \binom{n}{k}$ and its applications in number theory. II (Chinese)*, J. Nanjing Univ. Math. Biquarterly, **10**(1993), 105–118.
- [5] Z.H.Sun, *Congruences involving Bernoulli and Euler numbers*, J. Number Theory, **128**(2008), 280–312.
- [6] Z.-H. Sun and Z.-W. Sun, *Fibonacci numbers and Fermat's last theorem*, Acta Arith., **60**(1992), 371–388.
- [7] Z.-W. Sun, *On the sum $\sum_{k \equiv r \pmod{m}} \binom{n}{k}$ and related congruences*, Israel J. Math., **128**(2002), 135–156.
- [8] H. C. Williams, *A note on the Fibonacci quotient $F_{p-\varepsilon}/p$* , Canad. Math. Bull., **25**(1982), 366–370.
- [9] J. Wolstenholme, *On certain properties of prime numbers*, Quarterly Journal of Mathematics, **5**(1862), 35–39.

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA

E-mail address: haopan79@yahoo.com.cn